




Stuttering

Stuttering affects the flow or rhythm of verbal communication. People who stutter may repeat or prolong sounds and words in their speech.



What are the signs of stuttering?

People who stutter might show the following behaviours:

- Repetition of sounds, word parts, whole words, or phrases
- Sound prolongation, where sounds are stretched out in speech, e.g., sssssun
- Blocking of speech, where sounds get stuck, and no sound comes out for a period of time

These speech features may be accompanied by additional physical movements including:

- Facial tics (uncontrolled facial spasms such as rapid eye blinking or nose scrunching)
- Lip tremors
- Excessive eye blinking
- Hand flapping
- Tension in the face and upper body

How common is stuttering?

Stuttering is common, affecting approximately 1% of the population globally. Males are more likely to stutter than females. A large Australian study found that 8.5% of children will experience a period of stuttering by the age of 3 years, and 11.2% by the age of 4 years. While many children (up to 65%) recover naturally from stuttering, this is not always the case. Early intervention is recommended from a speech pathologist.

What causes stuttering?

Stuttering is not a language disorder or a psychological disorder. While little is known about the exact causes of stuttering, we know that both the brain and genes are involved.

Research studies have found:


- Stuttering can be associated with signal differences in the brain, nerves, and muscles
- Stuttering can run in families and has been linked to differences in genetic makeup

A smaller number of individuals may have stuttering associated with known genetic conditions such as Koolen-de Vries syndrome or Down syndrome.

Further research into the causes of stuttering may help to develop more targeted treatments in the future.

How is stuttering treated?

While many children naturally recover from stuttering, we are unable to predict who these children will be. It is therefore recommended that children who stutter receive treatment as early as possible. There are two broad types of treatment for stuttering that have been proven effective in clinical trials: The Lidcombe Program and Smooth Speech Treatment. A speech pathologist will help determine the treatment best suited to your child.



References

Boyce, J. O., et al. (2022). Self-reported impact of developmental stuttering across the lifespan. *Developmental medicine and child neurology*, 10.1111/dmcn.15211. Advance online publication. <https://doi.org/10.1111/dmcn.15211>

Brignell, A., et al. (2020). A systematic review of interventions for adults who stutter. *Journal of fluency disorders*, 64, 105766. <https://doi.org/10.1016/j.jfludis.2020.105766>

Brignell, A., et al. (2021). Interventions for children and adolescents who stutter: A systematic review, meta-analysis, and evidence map. *Journal of fluency disorders*, 70, 105843. <https://doi.org/10.1016/j.jfludis.2021.105843>

Genetics of Stuttering Study, retrieved from <https://www.geneticsofstutteringstudy.org.au/>

Kefalianos, E., et al. (2017). The History of Stuttering by 7 Years of Age: Follow-Up of a Prospective Community Cohort. *Journal of speech, language, and hearing research : JSLHR*, 60(10), 2828-2839. https://doi.org/10.1044/2017_JSLHR-S-16-0205

Reilly, S., et al. (2009). Predicting stuttering onset by the age of 3 years: a prospective, community cohort study. *Pediatrics*, 123(1), 270-277. <https://doi.org/10.1542/peds.2007-3219>

Reilly, S., et al. (2013). Natural history of stuttering to 4 years of age: a prospective community-based study. *Pediatrics*, 132(3), 460-467. <https://doi.org/10.1542/peds.2012-3067>